arXiv:1404.7690v2 [math.AT] 19 Mar 2015

TWISTED LEFSCHETZ NUMBERS OF INFRA-SOLVMANIFOLDS AND ALGEBRAIC GROUPS

HISASHI KASUYA

Abstract. Twisted Lefschetz numbers are extensions of the ordinary Lefschetz numbers for cohomologies with values in flat bundles. As a generalization of linearization formula for the ordinary Lefschetz number of a self-map of a nilmanifold, we show that a twisted Lefschetz number of any self-map of any infra-solvmanifold is equal to the determinant $\det(I - A)$ for some matrix A by using the cohomology of algebraic groups.

1. Introduction

Let M be a compact manifold and E be a flat vector bundle over M. We consider the cohomology $H^*(M, E)$ with values in E. Let $\varphi : M \to M$ be a smooth map and a morphism $\Xi : \varphi^* E \to E$ of flat bundle where $\varphi^* E$ is the pull-back of E. Consider the induced map $\Xi \circ \varphi^* : H^*(M, E) \to H^*(M, E)$. We define the twisted Lefschetz number

$$L(\varphi, E, \Xi) = \sum_i (-1)^i \mathrm{trace}(\Xi \circ \varphi^*)_{|H^i(M,E)}.$$

As the ordinary Lefschetz number, we can show the Lefschetz fixed point formula for the twisted Lefschetz number (see [1]). In general, computations of the twisted Lefschetz numbers are harder than computations of usual Lefschetz numbers. In this paper we give a simple formula for twisted Lefschetz numbers of maps of solvmanifolds and infra-solvmanifolds which is a generalization of "linearization formula" for ordinary Lefschetz numbers of maps of nilmanifolds.

In this paper solvmanifolds (resp. nilmanifolds) are compact homogeneous spaces of connected solvable (resp. nilpotent) Lie groups. We say that a solvmanifold (resp. nilmanifold) M has a discrete presentation (G, Γ) if there exists a simply connected solvable (resp. nilpotent) Lie group G with a cocompact discrete subgroup Γ so that M is diffeomorphic to G/Γ. It is known that every nilmanifold M has a discrete presentation (G, Γ) (see [9]).

Consider a nilmanifold G/Γ for a discrete presentation (G, Γ). For a smooth map $\varphi : G/\Gamma \to G/\Gamma$, since the fundamental group of G/Γ is Γ, we have the map $f : \Gamma \to \Gamma$ induced by φ. It is known that any homomorphism between cocompact discrete subgroups of simply connected nilpotent Lie groups can be extended to a Lie group homomorphism. Hence we have a extension $T : G \to G$ of f. In [12], Nomizu showed that the de Rham cohomology of a nilmanifold G/Γ for a discrete presentation (G, Γ) is isomorphic to the Lie algebra cohomology of the Lie algebra $\mathfrak{g}$ of G. Hence, considering the induced homomorphism $T_* : \mathfrak{g} \to \mathfrak{g}$, taking a matrix presentation A of $T_* : \mathfrak{g} \to \mathfrak{g}$, we can get the "linearization formula"

$$L(\varphi) = \det(I - A)$$

2010 *Mathematics Subject Classification.* 22E25; 20G20 ; 54H25; 55M20 .

Key words and phrases. cohomology of algebraic group, twisted lefschetz number, infra-solvmanifold.

where $L(\varphi)$ is the ordinary Lefschetz number.

We are interested in getting a "linearization formula" for solvmanifolds. But there are many difficulties:

- In general, a solvmanifold M do not have a discrete presentation (G, Γ).
- In general, for a simply connected solvable Lie group with a cocompact discrete subgroup Γ, a homomorphism $\Gamma \to \Gamma$ is not extended to a Lie group homomorphism $G \to G$.
- In general, the de Rham cohomology of a solvmanifold G/Γ for a discrete presentation (G, Γ) is not isomorphic to the Lie algebra cohomology of the Lie algebra $\mathfrak{g}$ of G.

Assuming some conditions, we can get a "linearization formula" (see for examples [10] and [8]). However on the usual Lefschetz number of any self-map of any solvmanifold, it seems to be difficult to get a "linearization formula".

In this paper we consider the more general setting. An infra-solvmanifold is a manifold of the form G/Δ, where G is a simply connected solvable Lie group, and Δ is a torsion-free subgroup of $\mathrm{Aut}(G) \ltimes G$ such that the closure of $h(\Delta)$ in $\mathrm{Aut}(G)$ is compact where $h : \mathrm{Aut}(G) \ltimes G \to \mathrm{Aut}(G)$ is the projection. An infra-solvmanifold is a generalization of a solvmanifold. The purpose of this paper is to give a "linearization formula" for any self-map of any infra-solvmanifold by using the twisted Lefschetz number and algebraic groups.

2. Rational cohomology of algebraic groups

Let $\mathcal{G}$ be a real algebraic group. We denote by $\mathcal{U}(\mathcal{G})$ the unipotent radical of a real algebraic group $\mathcal{G}$. It is known that the extension

$$1 \longrightarrow \mathcal{U}(\mathcal{G}) \longrightarrow \mathcal{G} \longrightarrow \mathcal{G}/\mathcal{U}(\mathcal{G}) \longrightarrow 1$$

splits [11]. We denote by $\mathbb{R}[\mathcal{G}]$ the coordinate ring of $\mathcal{G}$.

For a real algebraic group $\mathcal{G}$ and a rational $\mathcal{G}$-module V, we define the rational cohomology $H^*(\mathcal{G}, V) = \mathrm{Ext}^*_{\mathcal{G}}(\mathbb{R}, V)$ as [5] and [7]. We have

$$H^*(\mathcal{G}, V) = H^*(\mathcal{U}(\mathcal{G}), V)^{\mathcal{G}/\mathcal{U}(\mathcal{G})}$$

(see [5],[3]).

Consider the coordinate ring $\mathbb{R}[\mathcal{G}/\mathcal{U}(\mathcal{G})]$ of $\mathcal{G}/\mathcal{U}(\mathcal{G})$ as a $\mathcal{G}$-module. We have

$$H^*(\mathcal{G}, \mathbb{R}[\mathcal{G}/\mathcal{U}(\mathcal{G})]) = H^*(\mathcal{U}(\mathcal{G}), \mathbb{R})$$

(see [7, Part I. 4.7]). Since $\mathcal{G}/\mathcal{U}(\mathcal{G})$ is a reductive group, the module $\mathbb{R}[\mathcal{G}/\mathcal{U}(\mathcal{G}]$ is completely reductive. Hence for a sub-module $V \subset \mathbb{R}[\mathcal{G}/\mathcal{U}(\mathcal{G})]$, we have a projection $\mathbb{R}[\mathcal{G}/\mathcal{U}(\mathcal{G}] \to V$ and we have the inclusion $H^*(\mathcal{G}, V) \subset H^*(\mathcal{G}, \mathbb{R}[\mathcal{G}/\mathcal{U}(\mathcal{G})])$. Since $H^*(\mathcal{U}(\mathcal{G}), \mathbb{R})$ is finite-dimensional, we can take a finite dimensional sub-module $V \subset \mathbb{R}[\mathcal{G}/\mathcal{U}(\mathcal{G}]$ such that

$$H^*(\mathcal{G}, V) = H^*(\mathcal{G}, \mathbb{R}[\mathcal{G}/\mathcal{U}(\mathcal{G})]) = H^*(\mathcal{U}(\mathcal{G}), \mathbb{R}).$$

Consider the Lie algebra $\mathfrak{u}_{\mathcal{G}}$ of $\mathcal{U}(\mathcal{G})$ and the Lie algebra cohomology $H^*(\mathfrak{u}_{\mathcal{G}}, \mathbb{R})$. Then the functor from the category of rational $\mathcal{U}(\mathcal{G})$-modules to the category of $\mathfrak{u}_{\mathcal{G}}$-modules induces an isomorphism

$$H^*(\mathcal{U}(\mathcal{G}), \mathbb{R}) \cong H^*(\mathfrak{u}_{\mathcal{G}}, \mathbb{R})$$

(see [5]).

3. Group cohomology of torsion-free virtually polycyclic groups

In this section we consider representations of virtually polycyclic groups in algebraic groups.

Lemma 3.1. ([14, Lemma 4.36.]) *Let Γ be a torsion-free virtually polycyclic group. For a finite-dimensional representation $\rho : \Gamma \to GL_n(\mathbb{R})$, denote by $\mathcal{G}$ the Zariski-closure of $\rho(\Gamma)$ in $GL_n(\mathbb{R})$. Then we have* $\dim \mathcal{U}(\mathcal{G}) \leq \operatorname{rank} \Gamma$.

By this lemma, we consider the following definition.

Definition 3.2. Let Γ be a torsion-free virtually polycyclic group. For a real algebraic group $\mathcal{G}$, a representation $\rho : \Gamma \to \mathcal{G}$ is called full if the image $\rho(\Gamma)$ is Zariski-dense in $\mathcal{G}$ and we have $\dim \mathcal{U}(\mathcal{G}) = \operatorname{rank} \Gamma$.

We can compute the group cohomology of torsion-free virtually polycyclic groups by using rational cohomology of algebraic groups.

Theorem 3.3 ([6])**.** *Let Γ be a torsion-free virtually polycyclic group. We suppose that we have a real algebraic group $\mathcal{G}$ and an injection $\Gamma \subset \mathcal{G}$ which is a full representation. Let V be a rational $\mathcal{G}$-module. Then the inclusion $\Gamma \subset \mathcal{G}$ induces a cohomology isomorphism*

$$H^*(\mathcal{G}, V) \cong H^*(\Gamma, V).$$

For any torsion-free virtually polycyclic group, we have a good full representation.

Definition 3.4. We call a real algebraic group $\mathcal{A}_\Gamma$ a real algebraic hull of Γ if there exists an injective group homomorphism $\psi : \Gamma \to \mathcal{A}_\Gamma$ so that

- $\psi : \Gamma \to \mathcal{A}_\Gamma$ is a full representation.
- $Z_{\mathcal{A}_\Gamma}(\mathcal{U}(\mathcal{A}_\Gamma)) \subset \mathcal{U}(\mathcal{A}_\Gamma)$ where $Z_{\mathcal{A}_\Gamma}(\mathcal{U}(\mathcal{A}_\Gamma))$ is the centralizer of $\mathcal{U}(\mathcal{A}_\Gamma)$.

Theorem 3.5. ([2, Theorem A.1]) *There exists a real algebraic hull of Γ and a real algebraic hull of Γ is unique up to algebraic group isomorphism.*

4. Main result

Let M be an infra-solvmanifold and $\varphi : M \to M$ a smooth map. It is known that the fundamental group Γ of M is torsion-free virtually polycyclic. We consider the homomorphism $f : \Gamma \to \Gamma$ which is induced by $\varphi : M \to M$.

Take the real algebraic hull $\psi : \Gamma \to \mathcal{A}_\Gamma$ of Γ. Define $\Phi_f : \Gamma \to \mathcal{A}_\Gamma \times \mathcal{A}_\Gamma$ as $\Phi_f(\gamma) = (\psi(\gamma), \psi \circ f(\gamma))$. Let $\mathcal{G}_f$ be the real Zariski-closure of $\Phi_f(\Gamma)$ in $\mathcal{A}_\Gamma \times \mathcal{A}_\Gamma$. For the projection $p_1 : \mathcal{A}_\Gamma \times \mathcal{A}_\Gamma \ni (a, b) \mapsto (a) \in \mathcal{A}_\Gamma$, we consider the restriction $\Pi : \mathcal{G}_f \to \mathcal{A}_\Gamma$ of p_1 on $\mathcal{G}_f$. Then we have the commutative diagram

$$\begin{array}{ccc} \mathcal{G}_f & \xrightarrow{\Pi} & \mathcal{A}_\Gamma \\ & \nwarrow^{\Phi_f} & \uparrow{\scriptstyle \psi} \\ & & \Gamma \end{array} .$$

Since Φ_f and ψ have the Zariski-dense images, the map Π is surjective and hence $\Pi(\mathcal{U}(\mathcal{G}_f)) = \mathcal{U}(\mathcal{A}_\Gamma)$. By Lemma 3.1, $\dim \mathcal{U}(\mathcal{G}_f) = \dim \mathcal{U}(\mathcal{A}_\Gamma)$ and hence Π gives an isomorphism $\mathcal{U}(\mathcal{G}_f) \cong \mathcal{U}(\mathcal{A}_\Gamma)$. Thus $\Phi_f : \Gamma \to \mathcal{G}_f$ is a full representation.

For the projection $p_2 : \mathcal{A}_\Gamma \times \mathcal{A}_\Gamma \ni (a, b) \mapsto (b) \in \mathcal{A}_\Gamma$, we consider the restriction $F : \mathcal{G}_f \to \mathcal{A}_\Gamma$ of p_2 on $\mathcal{G}_f$. Then we have the commutative diagram

$$\begin{array}{ccc} \mathcal{G}_f & \xrightarrow{F} & \mathcal{A}_\Gamma \\ {\scriptstyle \Phi_f}\uparrow & & \uparrow{\scriptstyle \psi} \\ \Gamma & \xrightarrow{f} & \Gamma \end{array} .$$

Consider the restriction $F : \mathcal{U}(\mathcal{G}_f) \to \mathcal{U}(\mathcal{A}_\Gamma)$. Since we have an isomorphism $\Pi : \mathcal{U}(\mathcal{G}_f) \cong \mathcal{U}(\mathcal{A}_\Gamma)$, we obtain the linear map $S : \mathfrak{u}_{\mathcal{A}_\Gamma} \to \mathfrak{u}_{\mathcal{A}_\Gamma}$ which is the differential of the algebraic group homomorphism $F \circ \Pi^{-1} : \mathcal{U}_{\mathcal{A}_\Gamma} \to \mathcal{U}_{\mathcal{A}_\Gamma}$ where $\mathfrak{u}_{\mathcal{A}_\Gamma}$ is the Lie algebra of $\mathcal{U}_{\mathcal{A}_\Gamma}$.

Theorem 4.1. *We have a flat vector bundle E over M and a morphism $\Xi : \varphi^* E \to E$ of flat bundle such that*

$$L(\varphi, E, \Xi) = \det(I - A)$$

where A is a matrix presentation of $S : \mathfrak{u}_{\mathcal{A}_\Gamma} \to \mathfrak{u}_{\mathcal{A}_\Gamma}$.

Proof. By $H^*(\mathcal{A}_\Gamma, \mathbb{R}[\mathcal{A}_\Gamma/\mathcal{U}(\mathcal{A}_\Gamma)]) = H^*(\mathcal{U}(\mathcal{A}_\Gamma), \mathbb{R})$ and $H^*(\mathcal{G}_f, \mathbb{R}[\mathcal{G}_f/\mathcal{U}(\mathcal{G}_f)]) = H^*(\mathcal{U}(\mathcal{G}_f), \mathbb{R})$, the map $F^* : H^*(\mathcal{A}_\Gamma, \mathbb{R}[\mathcal{A}_\Gamma/\mathcal{U}(\mathcal{A}_\Gamma)]) \to H^*(\mathcal{G}_f, \mathbb{R}[\mathcal{G}_f/\mathcal{U}(\mathcal{G}_f)])$ is identified with $F^* : H^*(\mathcal{U}(\mathcal{A}_\Gamma), \mathbb{R}) \to H^*(\mathcal{U}(\mathcal{G}_f), \mathbb{R})$.

Take a finite dimensional sub-module $V \subset \mathbb{R}[\mathcal{A}_\Gamma/\mathcal{U}(\mathcal{A}_\Gamma)]$ such that

$$H^*(\mathcal{A}_\Gamma, V) = H^*(\mathcal{A}_\Gamma, \mathbb{R}[\mathcal{A}_\Gamma/\mathcal{U}(\mathcal{A}_\Gamma)]) = H^*(\mathcal{U}(\mathcal{A}_\Gamma), \mathbb{R})$$

as Section 2. We consider $\Pi^* V \subset \mathbb{R}[\mathcal{G}_f/\mathcal{U}(\mathcal{G}_f)]$. Since $\Pi : \mathcal{U}(\mathcal{G}_f) \to \mathcal{U}(\mathcal{A}_\Gamma)$ is an isomorphism, the map $\Pi^* : H^*(\mathcal{U}(\mathcal{A}_\Gamma), \mathbb{R}) \to H^*(\mathcal{U}(\mathcal{G}_f), \mathbb{R})$ is an isomorphism. We have the commutative diagram

$$\begin{array}{ccc} H^*(\mathcal{A}_\Gamma, V) & \xrightarrow{\Pi^*} & H^*(\mathcal{G}_f, \Pi^* V) \\ \| & & \downarrow \\ H^*(\mathcal{A}_\Gamma, \mathbb{R}[\mathcal{A}_\Gamma/\mathcal{U}(\mathcal{A}_\Gamma)]) & \xrightarrow{\Pi^*} & H^*(\mathcal{G}_f, \mathbb{R}[\mathcal{G}_f/\mathcal{U}(\mathcal{G}_f)]) \\ \| & & \| \\ H^*(\mathcal{U}(\mathcal{A}_\Gamma), \mathbb{R}) & \xrightarrow{\Pi^*(\cong)} & H^*(\mathcal{U}(\mathcal{G}_f), \mathbb{R}) \end{array} .$$

By this, we can say $H^*(\mathcal{G}_f, \Pi^* V) = H^*(\mathcal{G}_f, \mathbb{R}[\mathcal{G}_f/\mathcal{U}(\mathcal{G}_f)])$.

Since the module $\mathbb{R}[\mathcal{G}_f/\mathcal{U}(\mathcal{G}_f)]$ is completely reductive by the reductivity of $\mathcal{G}_f/\mathcal{U}(\mathcal{G}_f)$, we have the projection $\mathbb{R}[\mathcal{G}_f/\mathcal{U}(\mathcal{G}_f)] \to \Pi^* V$. Consider the restriction $\xi : F^* V \to \Pi^* V$ of this projection. The map $H^*(\mathcal{A}_\Gamma, V) \to H^*(\mathcal{G}_f, \Pi^* V)$ induced by F and ξ is identified with $F^* : H^*(\mathcal{U}(\mathcal{A}_\Gamma), \mathbb{R}) \to H^*(\mathcal{U}(\mathcal{G}_f), \mathbb{R})$. Consider V, $\Pi^* V$ and $F^* V$ as Γ-modules via Φ_f and ψ. Then we have $V = \Pi^* V$ as a Γ-module. By Theorem 3.3, the induced maps $\psi : H^*(\mathcal{A}_\Gamma, V) \to H^*(\Gamma, V)$, $\Phi_f : H^*(\mathcal{G}_f, F^* V) \to H^*(\Gamma, f^* V)$ and $\Phi_f : H^*(\mathcal{G}_f, \Pi^* V) \to H^*(\Gamma, V)$ are all isomorphisms.

By these arguments, we have the commutative diagram

$$\begin{array}{ccccccc}
H^*(\Gamma,V) & \xrightarrow{f^*} & H^*(\Gamma,f^*V) & \xrightarrow{\xi} & H^*(\Gamma,V) & = & H^*(\Gamma,V) \\
\uparrow{\scriptstyle \psi(\cong)} & & \uparrow{\scriptstyle \Phi_f(\cong)} & & \uparrow{\scriptstyle \Phi_f(\cong)} & & \uparrow{\scriptstyle \psi(\cong)} \\
H^*(\mathcal{A}_\Gamma,V) & \xrightarrow{F^*} & H^*(\mathcal{G}_f,F^*V) & \xrightarrow{\xi} & H^*(\mathcal{G}_f,\Pi^*V) & & H^*(\mathcal{A}_\Gamma,V) \\
\| & & & & \| & & \| \\
H^*(\mathcal{A}_\Gamma,\mathbb{R}[\mathcal{A}_\Gamma/\mathcal{U}(\mathcal{A}_\Gamma)]) & \xrightarrow{F^*} & & & H^*(\mathcal{G}_f,\mathbb{R}[\mathcal{G}_f/\mathcal{U}(\mathcal{G}_f)]) & & H^*(\mathcal{A}_\Gamma,\mathbb{R}[\mathcal{A}_\Gamma/\mathcal{U}(\mathcal{A}_\Gamma)]) \\
\| & & & & \| & & \| \\
H^*(\mathcal{U}(\mathcal{A}_\Gamma),\mathbb{R}) & \xrightarrow{F^*} & & & H^*(\mathcal{U}(\mathcal{G}_f),\mathbb{R}) & \xrightarrow{(\Pi^{-1})^*(\cong)} & H^*(\mathcal{U}(\mathcal{A}_\Gamma),\mathbb{R})
\end{array} .$$

As Section 2, we have

$$\begin{array}{ccc}
H^*(\mathcal{U}(\mathcal{A}_\Gamma),\mathbb{R}) & \xrightarrow{(F\circ\Pi^{-1})^*} & H^*(\mathcal{U}(\mathcal{A}_\Gamma),\mathbb{R}) \\
\downarrow{\scriptstyle \cong} & & \downarrow{\scriptstyle \cong} \\
H^*(\mathfrak{u}_{\mathcal{A}_\Gamma},\mathbb{R}) & \xrightarrow{(F\circ\Pi^{-1})^*} & H^*(\mathfrak{u}_{\mathcal{A}_\Gamma},\mathbb{R})
\end{array} .$$

Hence, taking a matrix presentation A of the Lie algebra homomorphism $S : \mathfrak{u}_{\mathcal{A}_\Gamma} \to \mathfrak{u}_{\mathcal{A}_\Gamma}$ induced by $F \circ \Pi^{-1}_{|\mathcal{U}(\mathcal{G}_f)} : \mathcal{U}_{\mathcal{A}_\Gamma} \to \mathcal{U}_{\mathcal{A}_\Gamma}$, we have $L(f,V,\xi) = \det(I-A)$ where $L(f,V,\xi) = \sum_i (-1)^i \mathrm{trace}(\xi \circ f^*)_{|H^i(\Gamma,V)}$.

Therefore, taking the flat bundle E which corresponds to V and the morphism $\Xi : \varphi^*E \to E$ of flat bundle which corresponds to ξ, we have

$$L(\varphi,E,\Xi) = L(f,V,\xi) = \det(I-A).$$

and hence we complete the proof of Theorem 4.1. □

Remark 1 (see [14], [13]). Suppose M is a nilmanifold. Then the fundamental group Γ of M is a finitely generated nilpotent group. In this case, the real algebraic hull $\mathcal{A}_\Gamma$ is unipotent (i. e. $\mathcal{A}_\Gamma = \mathcal{U}(\mathcal{A}_\Gamma)$) and hence we can take $V = \mathbb{R}$. Moreover $\mathcal{U}(\mathcal{A}_\Gamma)$ can be considered as a simply connected nilpotent Lie group which contains Γ as a cocompact discrete subgroup. Hence in nilmanifold case, Theorem 4.1 is the linearization formula for the ordinary Lefschetz number of a self-map of a nilmanifold as we explained in Introduction.

5. Application

We consider a solvmanifold G/Γ for a discrete presentation (G,Γ) with a smooth map $\varphi : G/\Gamma \to G/\Gamma$. Let $\mathfrak{g}$ be the Lie algebra of G. We suppose that the induced map $\varphi_* : \Gamma \to \Gamma$ can be extended to a Lie group homomorphism $T : G \to G$. Consider the induced map $T_* : \mathfrak{g} \to \mathfrak{g}$. If an isomorphism $H^*(G/\Gamma,\mathbb{R}) \cong H^*(\mathfrak{g},\mathbb{R})$ holds, then we have the equation

$$L(\varphi) = \det(I-B)$$

where B is a matrix presentation of $T_* : \mathfrak{g} \to \mathfrak{g}$. But in general an isomorphism $H^*(G/\Gamma,\mathbb{R}) \cong H^*(\mathfrak{g},\mathbb{R})$ do not hold. Thus the equation $L(\varphi) = \det(I-B)$ do not hold. Even if an isomorphism $H^*(G/\Gamma,\mathbb{R}) \cong H^*(\mathfrak{g})$ do not hold, it is desirable that the number $\det(I-B)$ gives some information for fixed points. As an application of Theorem 4.1, we have the following theorem.

Theorem 5.1. *Let G be a simply connected solvable Lie group with a cocompact discrete subgroup Γ and $\mathfrak{g}$ the Lie algebra of G. Let $\varphi : G/\Gamma \to G/\Gamma$ be a smooth map such that the induced map $\varphi_* : \Gamma \to \Gamma$ can be extended to a Lie group homomorphism $T : G \to G$. Take a matrix presentation B of the induced map $T_* : \mathfrak{g} \to \mathfrak{g}$. Then there exists a flat bundle E and a morphism $\Xi : \varphi^* E \to E$ of flat bundle such that*

$$L(\varphi, E, \Xi) = \det(I - B).$$

Proof. Let G be a simply connected solvable Lie group. For a finite dimensional representation $\rho : G \to GL_n(\mathbb{R})$, taking the Zariski-closure $\mathcal{G}$ of $\rho(G)$, we have $\dim \mathcal{U}(\mathcal{G}) \leq \dim G$ ([14, Lemma 4.36.]). As virtually polycyclic case, for a real algebraic group $\mathcal{G}$ and a representation $\rho : G \to \mathcal{G}$, we say that ρ is a full representation if the image $\rho(G)$ is Zariski-dense in $\mathcal{G}$ and we have $\dim \mathcal{U}(\mathcal{G}) = \dim G$.

We call a real algebraic group $\mathcal{A}_G$ a real algebraic hull of G if there exists an injective Lie group homomorphism $\psi' : G \to \mathcal{A}_G$ so that:

- $\psi' : G \to \mathcal{A}_G$ is a full representation.
- $Z_{\mathcal{A}_G}(\mathcal{U}(\mathcal{A}_G)) \subset \mathcal{U}(\mathcal{A}_G)$.

As similar to virtually polycyclic case, there exists an algebraic hull of G and an algebraic hull of G is unique up to algebraic group isomorphism (see [14, Proposition 4.40, Lemma 4.41]).

Take the real algebraic hull $\psi' : G \to \mathcal{A}_G$ of G. We suppose that G has a cocompact discrete subgroup Γ. Let $f : \Gamma \to \Gamma$ be a homomorphism which can be extended to a Lie group homomorphism $T : G \to G$. Define $\Phi_T : G \to \mathcal{A}_G \times \mathcal{A}_G$ as $\Phi_T(g) = (\psi'(g), \psi' \circ T(g))$. Let $\mathcal{G}_T$ be the real Zariski-closure of $\Phi_f(G)$ in $\mathcal{A}_G \times \mathcal{A}_G$. For the projection $p'_1 : \mathcal{A}_G \times \mathcal{A}_G \ni (a, b) \mapsto (a) \in \mathcal{A}_G$, we consider the restriction $\Pi' : \mathcal{G}_T \to \mathcal{A}_G$ of p'_1 on $\mathcal{G}_T$. Then we have the commutative diagram

$$\begin{array}{ccc} \mathcal{G}_T & \xrightarrow{\Pi'} & \mathcal{A}_G \\ & \nwarrow^{\Phi_T} & \uparrow{\scriptstyle \psi'} \\ & & G \end{array} .$$

For the projection $p'_2 : \mathcal{A}_G \times \mathcal{A}_G \ni (a, b) \mapsto (b) \in \mathcal{A}_G$, we consider the restriction $F' : \mathcal{G}_T \to \mathcal{A}_G$ of p_2 on $\mathcal{G}_T$. Then we have the commutative diagram

$$\begin{array}{ccc} \mathcal{G}_T & \xrightarrow{F'} & \mathcal{A}_G \\ {\scriptstyle \Phi_T}\uparrow & & \uparrow{\scriptstyle \psi'} \\ G & \xrightarrow{T} & G \end{array} .$$

As similar to Section 4, we can show that Π' gives an isomorphism $\mathcal{U}(\mathcal{G}_T) \cong \mathcal{U}(\mathcal{A}_G)$.

It is known that the real Zariski-closure of $\psi'(\Gamma)$ in $\mathcal{A}_G$ is the real algebraic hull $\mathcal{A}_\Gamma$ of a torsion-free polycyclic group Γ and $\mathcal{U}_{\mathcal{A}_\Gamma} = \mathcal{U}_{\mathcal{A}_G}$ (see [14, Proof of Theorem 4.34]). Hence, by restricting on Γ, the maps ψ', Φ_T, Π' and F' give the maps ψ, Φ_f, Π and F as in Section 4. In particular the map $F \circ \Pi^{-1} : \mathcal{U}_{\mathcal{A}_\Gamma} \to \mathcal{U}_{\mathcal{A}_\Gamma}$ is identified with the map $F \circ \Pi'^{-1} : \mathcal{U}(\mathcal{A}_G) \to \mathcal{U}(\mathcal{A}_G)$. Since the extension

$$1 \longrightarrow \mathcal{U}(\mathcal{A}_G) \longrightarrow \mathcal{A}_G \longrightarrow \mathcal{A}_G/\mathcal{U}(\mathcal{A}_G) \longrightarrow 1$$

splits, we have the projection map $q : \mathcal{A}_G \to \mathcal{U}(\mathcal{A}_G)$. Then it is known that the composition $Q = q \circ \psi' : G \to \mathcal{U}(\mathcal{A}_G)$ as a smooth map is a diffeomorphism (see [2,

Proposition 2.3]). Hence we have the commutative diagram

$$\begin{array}{ccc} \mathcal{U}(\mathcal{A}_\Gamma) & \xrightarrow{F\circ\Pi^{-1}} & \mathcal{U}(\mathcal{A}_\Gamma) \\ {\scriptstyle Q(\cong)}\uparrow & & \uparrow{\scriptstyle Q(\cong)} \\ G & \xrightarrow{T} & G \end{array} .$$

Hence the linear map $S : \mathfrak{u}_{\mathcal{A}_\Gamma} \to \mathfrak{u}_{\mathcal{A}_\Gamma}$ induced by $F \circ \Pi^{-1} : \mathcal{U}_{\mathcal{A}_\Gamma} \to \mathcal{U}_{\mathcal{A}_\Gamma}$ is identified with the linear map induced by $Q \circ T \circ Q^{-1}$. Hence taking a matrix presentation A of the Lie algebra homomorphism $S : \mathfrak{u}_{\mathcal{A}_\Gamma} \to \mathfrak{u}_{\mathcal{A}_\Gamma}$ and a matrix presentation B of the induced map $T_* : \mathfrak{g} \to \mathfrak{g}$, we have $\det(I - A) = \det(I - B)$.

In the setting of Theorem 5.1, taking the flat bundle E and the morphism $\Xi : \varphi^* E \to E$ of flat bundle as Section 4, we have $L(\varphi, E, \Xi) = \det(I - A)$. Hence we have $L(\varphi, E, \Xi) = \det(I - B)$.

Hence we complete the proof of Theorem 5.1.

□

References

[1] M. F. Atiyah, R. Bott, A Lefschetz fixed point formula for elliptic complexes. II. Applications. Ann. of Math. (2) **88** 1968 451–491.

[2] O. Baues, Infra-solvmanifolds and rigidity of subgroups in solvable linear algebraic groups. Topology **43** (2004), no. 4, 903–924.

[3] R. Hain, Remarks on non-abelian cohomology of proalgebraic groups. J. Algebraic Geom. **22** (2013), no. 3, 581–598.

[4] A. Hattori, Spectral sequence in the de Rham cohomology of fibre bundles. J. Fac. Sci. Univ. Tokyo Sect. I **8** 1960 289–331 (1960).

[5] G. Hochschild, Cohomology of algebraic linear groups. Illinois J. Math. **5** 1961 492–519.

[6] H. Kasuya, Central theorems for cohomologies of certain solvable groups. arXiv:1311.1310

[7] J. C. Jantzen, Representations of algebraic groups. Pure and Applied Mathematics, **131**. Academic Press, Inc., Boston, MA, 1987.

[8] J. Jezierski, W. Marzantowicz, Homotopy methods in topological fixed and periodic points theory. Topological Fixed Point Theory and Its Applications, **3**. Springer, Dordrecht, 2006.

[9] A. Malcev, On a class of homogeneous spaces. (Russian) Izvestiya Akad. Nauk. SSSR. Ser. Mat. **13**, (1949). 9–32.

[10] C. K. McCord, Nielsen numbers and Lefschetz numbers on solvmanifolds. Pacific J. Math. **147** (1991), no. 1, 153164.

[11] G. D. Mostow, Fully reducible subgroups of algebraic groups. Amer. J. Math. **78** (1956), 200–221.

[12] K. Nomizu, On the cohomology of compact homogeneous spaces of nilpotent Lie groups. Ann. of Math. (2) **59**, (1954). 531–538.

[13] A. L. Onishchik, E. B. Vinberg (Eds), Lie groups and Lie algebras II, Springer (2000).

[14] M.S. Raghnathan, Discrete subgroups of Lie Groups, Springer-Verlag, New York, 1972.

(Hisashi Kasuya) Department of Mathematics, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro, Tokyo 152-8551, Japan

E-mail address: kasuya@math.titech.ac.jp